

- Your partner(s) don't have to be involved.
- The patch lowers your chances of ovarian and endometrial cancer and fibroids.

Disadvantages of the patch

- You need a prescription.
- The patch only comes in one colour: light beige.
- You have to remember to remove and apply a new patch every week.
- If the patch comes off or starts to peel back, your risk of getting pregnant increases and you may need to use a back-up method of birth control for one week.
- The patch does not protect you from sexually transmitted infections (STIs).
- You may experience side effects.
- You may not be able to use the patch if you have certain health problems.
- If you have a personal and/or family history of breast cancer or you are HIV positive, talk to your clinician for more information.
- If you smoke and you use the patch, there is an increased risk of getting a blood clot.
- It can be expensive. However, PPT's Health Services sells the patch at a reduced cost.

For further information or to book an appointment:

For youth ages 13-29 - Planned Parenthood Toronto Health Services

Offers drop-in and scheduled appointments
Call 416-961-0113 or visit www.ppt.on.ca.

Teen Health Source

Offers anonymous and confidential sexual health information for teens by teens.
MSN chat TeenHealthSource@hotmail.com, email TeenHealthSource@ppt.on.ca,
call 416-961-3200 or visit www.TeenHealthSource.com.

For women - Bay Centre for Birth Control

Offers drop-in and scheduled appointments
Call 416-351-3700

Planned Parenthood Toronto is a United Way Member Agency and a Registered Charity,
No. 1190-94449-RR0001

Funding for Planned Parenthood Toronto has been provided by the Toronto Central Local
Health Integration Network.

REVISED JULY 2011

Birth Control Options Series



The Birth Control Patch

What is the birth control patch?

- The birth control patch (sometimes called the contraceptive patch or Evra) is a small, thin plastic patch containing 2 hormones (estrogen and progestin) that you wear on your skin.

How does the patch prevent pregnancy?

- In order to get pregnant, sperm must fertilize an egg that has been released. The patch prevents you from ovulating, which means your ovaries will not release an egg each month.
- The patch thickens the mucus on your cervix (the opening to your uterus). This makes it harder for sperm to travel into the uterus and fertilize an egg.
- The patch thins the lining of your uterus (the endometrium). This makes it harder for a fertilized egg to implant in the uterus and become a pregnancy.

How effective is the patch?

- The patch is 99% effective. This means that if 100 people used the patch correctly for one year, only one person would get pregnant.
- Because the patch may be used incorrectly, it is closer to 92% - 97% effective with typical use.
- If you use the patch incorrectly, your risk of getting pregnant increases.
- The patch may be less effective if you weigh more than 198 lbs.

How do you use the patch?

- The patch works on a 28-day (4 week) cycle and uses 3 patches per cycle.

From Choice, a World of Possibilities



planned
parenthood
TORONTO

- Week 1: apply the first patch on clean, dry, lotion-free skin.
- Week 2: remove first patch and apply second patch on the same day of the week that you applied your first patch.
- Week 3: remove second patch and apply third patch on the same day of the week that you applied your second patch.
- Week 4: remove third patch. Do NOT apply a new one. The next 7 days (week 4) will be “patch-free”. You can expect your period sometime during this week.
- At the end of that 7 day patch-free week, apply a new patch.
- Remember, you are protected from pregnancy during the patch-free week as long as you have been using the patch correctly and you apply your next patch on time.
- Some people use the patch continuously (no patch-free week) to avoid having their period, sometimes called “stacking”. If you are interested in doing this, speak to your clinician first.
- If you are late removing or applying your patch, consult the instructions in your patch package and/or call your clinician to see if you need to come in for an appointment and/or use a back-up method of birth control for 7 days.

How to start the patch

- If you decide, along with your clinician, that the patch is right for you, your clinician will write you a prescription. You can purchase your patches at a pharmacy or at PPT’s Health Services for a reduced cost.
- If you start the patch within the first 5 days of getting your period, you are protected from pregnancy right away.
- If you start the patch 6 or more days after getting your period, you are not protected from pregnancy until you have been using the patch for a full week. To avoid pregnancy during this time, you should use a back-up method of birth control like condoms or spermicides.
- A clinician may recommend that you use a back-up method of birth control for a longer period of time when you start the patch.
- Some people like to start the patch on the first Sunday following the start of their period, whether they are still bleeding or not. This will likely keep you from getting future periods on the weekend.

Where can I wear the patch?

- You can wear the patch on your butt, back, the outside of your arm or on your lower abdomen (stomach). Do not place the patch on or near your breasts.
- You can shower, swim, exercise and do regular activities while wearing the patch. In 98% of cases, the patch does not fall off.

What are the side effects of the patch?

- Some patch users will experience minor side effects such as nausea, sore breasts, moodiness and/or spotting (a little bit of bleeding that is lighter than your period). These usually go away within the first 3 months of using the patch.
- Some patch users will experience skin irritation on the spot that you wear your patch. Every time you change the patch, switch sides or apply it to a different site to avoid skin irritation.
- If after 3 months you are still experiencing side effects or your side effects are severe, you may want to try a different method of birth control.
- There is a rare risk of getting blood clots, or having a heart attack or stroke while using the patch. Smoking, obesity, and other health conditions increase this risk. Speak to your clinician for more information.
- Signs of a blood clot include: blurred or loss of vision, chest pain or difficulty breathing, migraine headaches, severe abdominal cramps, or severe pain in the leg. If you experience any of these symptoms, get medical attention right away.

Advantages of the patch

- If you use the patch correctly, your chances of getting pregnant are very low.
- You don’t have to do something every day and you are less likely to have side effects than you are with the pill.
- Your period will likely become shorter, lighter, less painful, and more regular.
- The patch can improve acne and increase bone strength.
- You don’t have to do anything before or after you have sex.
- It doesn’t affect your ability to get pregnant in the future.